

1. Report No. FAA-AM-78-18		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle EXPERIMENTAL ATTEMPTS TO EVOKE A DIFFERENTIAL RESPONSE TO DIFFERENT STRESSORS				5. Report Date	
				6. Performing Organization Code	
7. Author(s) C. E. Melton, J. M. McKenzie, J. T. Saldivar, and S. M. Wicks				8. Performing Organization Report No.	
9. Performing Organization Name and Address FAA Civil Aeromedical Institute P. O. Box 25082 Oklahoma City, Oklahoma 73125				10. Work Unit No. (TRAIS)	
				11. Contract or Grant No.	
12. Sponsoring Agency Name and Address Office of Aviation Medicine Federal Aviation Administration 800 Independence Avenue, S.W. Washington, D.C. 20591				13. Type of Report and Period Covered	
				14. Sponsoring Agency Code	
15. Supplementary Notes Work was performed under Task AM-C-77-PHY-103.					
16. Abstract Ten paid male subjects each worked at a physical task with no competitive element (treadmill) and a competitive task ("Pong") with minimal physical activity. There were three work periods, each 50 min long. Ten minutes were allowed for rest and urine collection after each work period. The experimental period lasted 3 h. Urine was analyzed for 17-ketogenic steroids (17-KGS), epinephrine (E), and norepinephrine (NE). Heart rates were derived from ambulatory electrocardiograms. There were no statistically significant differences in excretion of urinary metabolites during corresponding episodes of the two tasks. Heart rates were significantly higher during treadmill work than during Pong playing. Rest-to-work differences show that the increment in E excretion is significantly greater during the Pong task than during the treadmill task. Rest-to-work differences in excretion of 17-KGS and NE are not significant. The rest-to-work increase in heart rate is significant for treadmill, but not for Pong. The increase in epinephrine excretion strengthens the conclusion drawn from field experiments that this measurement is the best indicator of the intensity of air traffic control work <u>per se</u> .					
17. Key Words Stress Urinary Metabolites Air Traffic Controllers			18. Distribution Statement Document is available to the public through the National Technical Information Service, Springfield, Virginia 22161.		
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 5	22. Price

to score 15 points wins the game. One of the researchers acted as opponent for all the subjects; she was an expert at the game and was rarely beaten.

On arrival at the laboratory, subjects were requested to void urine and discard it. They then had electrocardiographic electrodes attached to their chests, were given 250 ml of water to drink, and were asked to rest in the supine position on a cot for 50 min. At the end of the rest period, subjects collected a urine specimen and began the first work episode.

The electrocardiogram was recorded on an Avionics Electrocardiocorder for continuous registration of heart rate. Urine specimens were collected in a 500-ml graduate cylinder, the volume was recorded, and aliquots were taken for analysis of 17-ketogenic steroids (17-KGS), epinephrine (E), and norepinephrine (NE). Aliquots were kept frozen in a freezer until analyzed. Urinary stress hormone values are expressed as total weights of the substances excreted during each 50-min episode.

III. Results.

The results of urine and heart rate analyses are shown in Tables 1 and 2. There are no statistically significant differences in levels of urinary metabolite excretion for corresponding episodes of the two tasks. Heart rates are significantly higher for the treadmill than for the Pong task (Table 1). Rest-to-work differences show that the increment in E excretion is significantly greater during the Pong task than during the treadmill task. Rest-to-work differences in excretion of 17-KGS and NE are not significant for either task. The rest-to-work increase in heart rate is significant for the treadmill but not for the Pong task (Table 2).

IV. Discussion.

Field experiments have shown that epinephrine excretion is significantly related to traffic count and to radio transmission time (1). The data have strongly suggested that adrenal steroid excretion is related to chronic stressors such as labor-management difficulties (2,3) and that norepinephrine excretion is related to physical activity (4). These experiments strengthen the interpretation that epinephrine excretion is related to mental tasks (5) such as air traffic control and not to physical tasks and therefore is the best single indicator of response to air traffic control work per se.

TABLE 1. Comparison of Excretion Values and Heart Rates for Pong and Treadmill Tasks*

Task	Total Amounts of Hormones Excreted			Heart Rate (Beats Per Minute)
	17-KGS mg	E ng	NE ng	
Rest (Pong)	0.70	1,237	3,603	64
Rest (T-Mill)	0.67	1,214	4,274	64
P	NS**	NS	NS	NS
Pong 1	0.70	1,619	3,809	73
T-Mill 1	0.59	1,741	4,384	101
P	NS	NS	NS	0.05
Pong 2	0.62	1,720	3,379	73
T-Mill 2	0.67	1,463	3,813	100
P	NS	NS	NS	0.05
Pong 3	0.59	1,750	3,833	70
T-Mill 3	0.58	1,491	3,581	98
P	NS	NS	NS	0.01

* Group Averages

** T-test

TABLE 2. Statistical Significance of Rest-To-Work Differences for the Various Measurements*

TASK	Level of Significance of Difference Between Rest and Task (P**)			HEART RATE
	17-KGS	E	NE	
Pong 1	NS	0.01	NS	NS
Pong 2	NS	0.01	NS	NS
Pong 3	NS	0.05	NS	NS
T-Mill 1	NS	NS	NS	0.01
T-Mill 2	NS	NS	NS	0.01
T-Mill 3	NS	NS	NS	0.01

* See Table 1 for actual values.

** Paired t-test

REFERENCES

1. Melton, C. E., J. M. McKenzie, J. T. Saldivar, and S. M. Hoffmann: Comparison of Opa Locka Tower With Other ATC Facilities by Means of a Stress Index, FAA Office of Aviation Medicine Report No. AM-74-11, 1974.
2. Melton, C. E., J. M. McKenzie, J. T. Saldivar, and Marlene Hoffmann: Studies on Stress in Aviation Personnel: Analysis and Presentation of Data Derived From a Battery of Measurements, Advisory Group for Aerospace Research and Development No. CPP-180, 1975.
3. Kagan, A. R., and L. Levi: Health and Environment - Psychosocial Stimuli, A Review. Reports from the Laboratory for Clinical Stress Research No. 27, December 1971.
4. Adkins, J. R., T. I. Miller, T. Carter, and J. D. Hardy: The Hormonal Mediation of Neurovascular Reflex Adjustments: Catecholamine Response to Postural Changes in Man, AMER. J. SURG., 27:210, 1961.
5. Pátkai, P.: Catecholamine Excretion in Pleasant and Unpleasant Situations, ACTA PSYCHOLOGICA, 35:352-363, 1971.

