

FAA-AM-89-2

A/AM-89/2

Doc
FAA
AM
89
02

Aviation Medicine
ton, D.C. 20591

Prevalence of Disease Among Active Civil Airmen



Charles F. Booze, Jr., Ph.D.

Civil Aeromedical Institute
Office of Aviation Medicine
Federal Aviation Administration
Oklahoma City, OK 73125

October 1988

Final Report

This document is available to the public
through the National Technical Information
Service, Springfield, Virginia 22161



U.S. Department of Transportation
Federal Aviation Administration

Doc
FAA
AM
89/02

E.R.A.U. LIBRARY

1. Report No. DOT/FAA/AM-89/2		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle PREVALENCE OF DISEASE AMONG ACTIVE CIVIL AIRMEN				5. Report Date OCTOBER 1988	
				6. Performing Organization Code	
				8. Performing Organization Report No.	
7. Author(s) Charles F. Booze, Jr., Ph.D.				10. Work Unit No. (TRAIS)	
9. Performing Organization Name and Address FAA Civil Aeromedical Institute P.O. Box 25082 Oklahoma City, Oklahoma 73125				11. Contract or Grant No.	
				13. Type of Report and Period Covered	
				14. Sponsoring Agency Code AAM-330	
12. Sponsoring Agency Name and Address Office of Aviation Medicine Federal Aviation Administration 800 Independence Avenue, S.W. Washington, D.C. 20591				15. Supplementary Notes	
16. Abstract It has been the policy of the Federal Aviation Administration to medically certify, for a variety of flying privileges, individuals who have a medical deficiency or disease, provided it can be determined that such action does not compromise air safety. This descriptive epidemiologic study presents the point prevalence of pathology among 685,552 active airmen as of January 1, 1988, by major body system. The study also presents other selected pathologies of interest within the major body system. Cardiovascular, eye, and abdominal pathologies represent the most prevalent medical conditions among active airmen (7.5%, 6.4% and 6.3% respectively). Hypertension is the most frequently occurring cardiovascular condition. Prevalence of pathology among active civil airmen, while still considerably less than among the general population, is increasing due to recent emphasis on relaxation of regulatory requirements when consistent with safety.					
17. Key Words Prevalence Disease Active airmen			18. Distribution Statement Document is available to the public through the National Technical Information Service, Springfield, VA 22161.		
19. Security Classif. (of this report) UNCLASSIFIED		20. Security Classif. (of this page) UNCLASSIFIED		21. No. of Pages 10	22. Price

NOTICE

This document is disseminated under the sponsorship of the U.S. Department of Transportation in the interest of information exchange. The United States Government assumes no liability for the contents or use thereof.

PREVALENCE OF DISEASE AMONG
ACTIVE CIVIL AIRMEN

INTRODUCTION

The Federal Aviation Administration (FAA) and its predecessors have been charged with the responsibility for medical certification of all United States and some international civil airmen since 1927. Each airman must hold a current medical certificate of the appropriate class to validate any pilot certificates he or she may possess. As of January 1, 1988, 685,552 active airmen were medically certified. Federal Aviation Regulations require that physical examinations be performed at 6-month intervals for air transport pilots, annually for other commercial pilots, and at 2-year intervals for private pilots.

The Aeromedical Certification Branch of the Civil Aeromedical Institute (CAMI), located in Oklahoma City, Oklahoma, is the central screening facility and repository within the FAA for collection, processing, adjudication, investigation, and analyses of medical data generated by the aeromedical certification and related regulatory programs.

Medical certification criteria have changed dramatically in favor of the airman during recent years as a result of the evolution of aviation medicine and increased efforts in the area of aeromedical research. A primary function of CAMI is to identify and provide substantive research results to verify or incriminate current medical criteria in the furtherance of aviation safety and better service to the airman. It has been the policy of the FAA to medically certify for a variety of flying privileges, individuals who also have medical deficiency or disease, provided it can be determined that such action does not compromise air safety.

This descriptive epidemiologic study presents the period prevalence of pathology among active airmen for years 1986 and 1987 by age, class of medical certificate, major body system, and other selected pathologies of interest within the major body systems. The study updates previous studies of a similar nature with respect to current prevalence of disease among airmen.

METHODS

Physical examinations to detect medical conditions which could incapacitate a pilot or otherwise adversely affect pilot performance are given by some 6,799 designated aviation medical examiners (AME's), most of whom are physicians in private practice. Military applicants receive their examinations at 410 designated military facilities. Reports of these examinations from throughout the world are forwarded to the Aeromedical Certification Branch in Oklahoma City.

Federal Aviation Regulations, Part 67, specify that a medical certificate will be denied if an applicant has an established medical history or clinical diagnosis of any of the following conditions:

1. A personality disorder that is severe enough to have repeatedly manifested itself by overt acts.
2. A psychosis.
3. Alcoholism, unless there is established clinical evidence, satisfactory to the Federal Air Surgeon, of recovery, including sustained total abstinence from alcohol for not less than the preceding 2 years. "Alcoholism" means a condition in which a person's intake of alcohol is great enough to damage physical health or personal or social functioning, or when alcohol has become a prerequisite to normal functioning.
4. Drug dependence.
5. Epilepsy.
6. A disturbance of consciousness without satisfactory medical explanation of the cause.
7. Myocardial infarction.
8. Angina pectoris.
9. Coronary heart disease that has required treatment or, if untreated, that has been symptomatic or clinically significant.
10. Diabetes mellitus, requiring insulin or other hypoglycemic drug for control.

However, certification is possible despite the existence of one of the above disqualifying medical conditions if special issuance is granted after extensive medical review by FAA and consultant specialists. The primary considerations in such cases are history, findings, prognosis, and potential risk of sudden or insidious incapacitation. It is appropriate to note, however, that when airmen with disqualifying conditions are issued medical certificates it is with special medical and operational restrictions that allow for control of risk.

Automated medical record files maintained by the Aeromedical Certification Branch provided the source data for the study. The files utilized contained only the most recent application for medical certification from an airman within the past 3 years. Pathology detected by previous medical examinations is brought forward to the current automated medical file in order to maintain as complete a history as possible. While this file contains records for a total of 3 years, a medically certified airman is considered "active" for a maximum of 24 calendar months following his or her most recent FAA medical examination; i.e., regardless of the class of medical certificate originally issued, it is valid for third-class airman purposes for a period of time up to 24 calendar months unless otherwise limited or recalled by the FAA. This study includes airmen certified during 1986 and 1987.

Prevalence data were automatically produced from the automated files described above. Prevalence data in this population are expected to be somewhat conservative since problems have been experienced in the acquisition of complete and accurate histories from pilots with a job, hobby, or aircraft investment to protect.

Prevalence data are obviously a function of incidence and duration of diseases; therefore, one would not expect the more immediately lethal diseases to be represented to the same extent as less serious diseases of longer duration. Additionally, airmen with more serious health problems are likely to be denied medical certification and would not become eligible for study. These data do, however, indicate the relative importance of various diseases among airmen, even though conservative in some instances, and provide insight concerning potential health problem interaction with the aviation environment. Since general population prevalence data are limited with respect to some of the diseases reported on in this study, some further benefit is expected in that regard.

RESULTS AND DISCUSSION

Diseases of the cardiovascular system are the most frequently observed diseases among active airmen. As expected, hypertension is the greatest contributor to the total prevalence rate. However, as may be seen in Table I, the prevalence of hypertension is only a fraction of that which would be expected in an unscreened population. The low prevalence of hypertension in this population is due to higher than usual upper limit cutoffs for assignment of a hypertensive diagnosis among airmen (170/100 for applicants for second- and third-class certification, and only slightly more stringent for class one applicants). Medications acceptable to the FAA for treatment of hypertension include all diuretics, all Food and Drug Administration approved beta-adrenergic blockers including combination agents, ACE inhibitors, calcium slow channel blocking agents, and combinations thereof.

Heart murmurs including functional or physiologic murmurs, account for over 16 percent of all heart conditions observed to exist among active airmen.

Reference to Tables I and II indicates that eye diseases are the next most prevalent medical problems among active airmen (excluding lens correction). While representing a potential threat to air safety if not recognized and treated by adequate correction, most eye pathology is relatively innocuous. Of these 685,552 active airmen as of January 1, 1988 some 48 percent (326,757) required lens correction for some visual deficiency. Of this total, 21,650 are known contact lens wearers. Additionally, 3,984 airmen are certified to fly with blindness or absence of an eye (which includes visual acuity worse than 20/200 uncorrected in either eye). Serious incapacitation during flight is possible should corrective lenses become dislocated and temporarily unavailable during a critical phase or flight. Therefore, considerable research effort has been and will continue to be devoted to this important aspect of potentially health-related pilot dysfunction.

Prevalence of abdominal diseases is third in total body system importance. Specific diseases making major contributions include current hernias, history of kidney stones, and uncomplicated ulcers.

Major body system prevalence rates for neuropsychiatric and bone and joint diseases are next highest at 21.2 and 16.6 per 1,000 airmen respectively. Within these systems no single disease entity makes a contribution significant enough to deserve separate mention. Both major body systems contain 30 to 40 distinct disease categories, with each making minor contributions to the total for the major body system.

TABLE I. TEN MOST PREVALENT CONDITIONS AMONG ACTIVE AIRMEN
 BY AGE
 (Rate per 1,000 Airmen)

Disease	Age (Years)					Total
	< 30	30-39	40-49	50-59	59 >	
Hypertension	4.5	16.9	36.9	70.0	98.2	31.6
Wears Contact lens	28.7	42.1	33.6	18.4	15.0	31.2
Kidney Stones	4.3	15.5	25.8	31.6	30.7	18.2
Hernia	5.7	10.6	19.9	33.1	50.7	17.5
Hay Fever	11.7	16.6	17.7	14.8	12.2	15.0
Heart Murmur	14.0	12.4	10.1	10.9	17.4	12.4
History of Neoplasm	1.5	4.0	8.6	17.5	38.5	8.8
Asthma	6.9	6.9	6.8	5.9	5.4	6.6
Uncomplicated Ulcer	2.9	5.5	8.2	10.0	11.0	6.5
Arthritis	0.4	2.5	7.1	14.0	23.7	6.1

TABLE II. PATHOLOGY PREVALENCE AMONG ACTIVE AIRMEN
 BY AGE AND MAJOR BODY SYSTEMS
 (Rate per 1,000 Airmen)

Body System	Age (Years)					Total	Rank
	< 30	30-39	40-49	50-59	59 >		
Eye	40.8	66.3	67.3	70.0	116.9	63.8	2
ENT	2.4	5.2	11.2	17.8	36.6	9.9	6
Respiratory	1.1	2.4	3.5	5.8	8.7	3.3	7
Cardiovascular	27.1	45.6	80.8	143.5	221.8	75.3	1
Abdominal (GI, GU, etc.)	20.7	46.7	76.9	110.6	153.1	63.4	3
Neuropsychiatric	13.0	20.2	27.0	27.8	24.4	21.2	4
Bones and Joints	5.0	10.4	18.3	30.8	50.9	16.6	5
Muscles	0.5	1.1	2.2	3.0	5.4	1.8	8
Miscellaneous	32.6	44.1	59.3	74.8	89.8	52.2	

The "miscellaneous" category of disease includes endocrinopathies, skin diseases, drug usage by type, allergies, tropical diseases, and general systemic conditions. Nearly 30 percent of all disease observed in this category was "hay fever." Other major contributors to this category included antihistamine drug usage, asthma, diabetes controlled by diet, and hypothyroidism.

The ten most prevalent diseases observed among active airmen are presented in Table I. Diseases found to increase with age include hypertension, kidney stones, hernia, history of neoplasm, uncomplicated ulcer and arthritis. Diseases demonstrating a trend of decrease with age included contact lens use, hay fever, and asthma. Heart murmurs were most common among young and older ages with lower prevalence among the middle ages.

These age/disease findings are generally consistent with expectation except for the decreasing prevalence of hay fever with age, which may be due to underreporting.

Overall, disease is slightly more prevalent among third-class (general aviation) airmen than among class one and two airmen (professional pilots) (see Figure 1). Since the professional pilot undergoes a more thorough and rigid examination and is subjected to stricter standards by the FAA as well as employers, one must assume that these results reflect the effects of more thorough early screening out of problem medical cases which results in less subsequent disease among the remaining group. Possibly medical history (which would lead to disease detection) is being masked to a greater extent among the professional-category pilots. Certainly the incentive is great, both monetary and otherwise, for the latter hypothesis.

The prevalence findings of this study and previous research efforts involving the airman population or segments thereof are generally similar with respect to major body systems presenting the highest prevalence rates; i.e., cardiovascular, eye, and abdominal (1-3). However, as seen in Figure 2, prevalence rates are higher than those in previous studies for most major body systems partially due to the fact that prevalence, by definition, is an accumulation of disease and would be greater in a given population at successive measurements for nonlethal diseases. The second consideration with regard to increased prevalence rates surely has to do with the fact that medical requirements have been cautiously relaxed over recent years to allow airmen to continue flying with various medical problems not previously acceptable. A possible additional factor having to do with the increase in prevalence seen in this study has to do with a rather dramatic decrease in the airman population. It seems likely that the residual population are more "hard core" aviators, many of whom may have accumulated pathology over the years and are thus still represented in the numerator while the denominator has decreased significantly..

Large and statistically significant increases in age-specific cardiovascular pathology were observed in comparing this study with the 1984 study. This difference is largely attributable to an increased number of hypertensives in the population resulting from a relaxation of allowable medications and combinations, and more liberal certification of airmen with cardiovascular disease during recent years.

Sizable increases in age-specific rates for abdominal pathologies were also observed in this study over the previous study. No single explanation, however, accounts for this latter observation other than that speculated earlier.





Figure 1. Disease Prevalence Among Active Airmen by Class of Medical Certificate, 1988.

RATES PER 1,000 AIRMEN

LEGEND





Figure 2. Comparison of Disease Prevalence Among Airmen by Body System, 1988 Verses 1984

SUMMARY

Health findings in the active airman population are generally consistent with those that might be expected in the general U.S. population, except scaled to reflect lower prevalence due to prescreening and problems associated with accurate reporting of symptoms and history by pilots.

Cardiovascular diseases are highest in frequency of occurrence among active airmen. As expected, hypertension is the greatest contributor to cardiovascular disease prevalence.

Eye disease is the next most common problem among pilots with some 48 percent of all airmen requiring lens correction for some visual deficiency.

Abdominal disease prevalence is third in total body system importance. Major contributors to the category include current hernia conditions, history of kidney stones, and uncomplicated ulcers.

Overall, disease prevalence is greater among currently certified airmen than among previous groups studied. This increase in prevalence is probably a reflection of more liberal standards more than any other single factor. Through the years the trend has been toward cautious relaxation of medical standards where possible when not resulting in a compromise of air safety.

The difference in prevalence of disease among general aviation pilots compared to professional pilots is much less than in previous studies and certainly reflects a more liberal certification policy by FAA in recent years. However, in view of the trend away from airline medical departments, and the preventive medical benefits of such departments, possibly we are seeing the result of diminished preventive efforts.

REFERENCES

1. Booze, C., 1973, Prevalence and incidence of disease among airmen medically certified during 1965. FAA Office of Aviation Medicine Report No. AM-73-8.
2. Booze, C., 1979. Morbidity experience of air traffic control personnel - 1967-77. Aviation, Space, and Environmental Medicine, 59(1):1-8.
3. Booze, C., 1981. Prevalence of selected pathology among currently certified active airmen. FAA Office of Aviation Medicine Report No. AM-81-9