## AD-A268 661

DOT/FAA/AM-93/10

Office of Aviation Medicine Washington, D.C. 20591



## Head and Face Anthropometry of Adult U.S. Civilians


Joseph W. Young

Civil Aeromedical Institute Federal Aviation Administration Oklahoma City, Oklahoma 73125

July 1993


Final Report

This document is available to the public through the National Technical Information Service, Springfield, Virginia 22161.





U.S. Department of Transportation Federal Aviation Administration



## NOTICE

This document is disseminated under the sponsorship of the U.S. Department of Transportation in the interest of information exchange. The United States Government assumes no liability for the contents or use thereof.

1. Report No.	2. Government Accession No.	<del></del>	3. Recipient's Catalog No.		
DOT/FAA/AM-93/10					
4. Title and Subtitle			5. Report Date		
Head and Face Anthropometi	ry of Adult U.S. Citize	ens	July 1993		
			6. Performing Organization	Code	
7. Author(s)			8. Performing Organization	Report No.	
Joseph W. Young					
Performing Organization Name and Address			10. Work Unit No. (TRAIS)		
FAA Civil Aeromedical Instit	tute				
P.O. Box 25082					
Oklahoma City, OK 73125			11. Contract or Grant No.		
12. Sponsoring Agency name and Address			13. Type of Report and Per	iod Covered	
Office of Aviation Medicine Federal Aviation Administrati	ion				
800 Independence Avenue, S.W.					
Washington, DC 20591		14. Sponsoring Agency Code			
15. Supplemental Notes					
Work was performed under Ta	ask AM-B-92-PRS-87 1	6.			
40 Made at					
This report presents a total of	17 traditional and 5 nev	v head and fa	cial dimensions from	a random.	
composite U.S. female and ma					
measurements, identified to de					
influence on equipment design adult subjects in the age range of					
subjects to other ethnic origin	subjects. The data sum	mary format	is specifically intende	ed for basic design	
concept use. These dimensions					
in the development of protective of anatomical and anthropome					
terms, specific to descriptive la				<u> </u>	
anthropometrical dimensions.					
17. Key Words		18. Distribution St			
Anatomy, Anthropometry, Anatomical		Document is available to the public through the			
Characteristics, Head and Face	: Anthropometry		echnical Information Virginia 22161.	Service,	
19. Security Classif. (of this report)	20. Security Classif. (of this page)	<u> </u>	21. No. of Pages	22 Price	
Unclassified	Unclas	sified	44		

Form DOT F 1700.7 (8-72)


Reproduction of completed page authorized

## **FOREWORD**

This report presents a current CAMI data base of adult head and face anthropometry that will typically identify a generalized morphological diversity of the U. S. population. The initial data format and data for this CAMI data base was generated, in support of a North American Aviation, Inc. research and development program, to provide special design criteria for the protective helmets, protective breathing devices, and pressurized flight gear for advanced weapon systems. These data, which form a part of the CAMI/1992 data base, represent both traditional and new unique measurements.

The largest part of the CAMI data sample was obtained in support of long range research and development programs at the Civil Aeromedical Institute, Federal Aviation Administration, Department of Transportation, and related protective equipment industries. These data support past and ongoing CAMI joint research and evaluation programs with design criteria for face related protective equipment and identify the ranges of face morphology of test subjects for comparison with target populations.

## DITC QUALITY INSPECTED 3



## TABLE OF CONTENTS

	N1
ANTHROPOM	ETRIC DATA BASES
STATISTICAL D	ESCRIPTION
<b>DEFINITIONS C</b>	OF ANATOMICAL LANDMARKS AND PLANES
<b>DEFINITIONS C</b>	OF ANTHROPOMETRICAL MEASUREM 12
REFERENCES	14
APPENDIX A	ANTHROPOMETRY THESAURUS A-1
APPENDIX B	STATISTICAL SUMMARY TABLES
	·
	LIST OF FIGURES
FIGURE 1.	COMPARATIVE HEAD BREADTH PERCENTILES
FIGURE 2.	COMPARATIVE HEAD LENGTH PERCENTILES
FIGURE 3.	COMPARATIVE FACE BREADTH PERCENTILES4
FIGURE 4.	COMPARATIVE FACE LENGTH PERCENTILES4
FIGURE 5.	COMPARATIVE NOSE BREADTH PERCENTILES
FIGURE 6.	COMPARATIVE HEAD LENGTH PERCENTILES
FIGURE 7.	FACE & HEAD LANDMARKS (ANTERIOR VIEW)8
FIGURE 8.	FACE & HEAD LANDMARKS (LATERAL VIEW)8
FIGURE 9.	FACE & HEAD LENGTH MEASUREMENTS (LATERAL VIEW)9
FIGURE 10.	FACE BREADTH MEASUREMENTS (ANTERIOR VIEW)9
FIGURE 11.	FACE & HEAD BREADTH MEASUREMENTS & REFERENCE PLANE LINES FOR
	HEIGHT MEASUREMENTS (ANTERIOR VIEW)10
FIGURE 12.	NASAL BRIDGE HEIGHT MEASUREMENTS FROM REFERENCE PLANE LINES
	(LATERAL VIEW)11
FIGURE 13.	PRONASALE HEIGHT MEASUREMENTS FROM REFERENCE PLANE LINES
	(LATERAL VIEW)11
	LIST OF TABLES
REFERENC	
TABLE 1.	REFERENCE NUMBERS OF ANTHROPOMETRIC MEASUREMENTS
STATISTIC	AL SUMMARY TABLES
TABLE 2.	HEAD CIRCUMFERENCEB-1
TABLE 3.	HEAD BREADTH
TABLE 4.	HEAD LENGTH B-3
TABLE 5.	BIORBITAL BREADTH
TABLE 6.	BIECTOCANTHUS BREADTHB-5
TABLE 7.	BIPUPIL BREADTHB-6

## STATISTICAL SUMMARY TABLES (Continued)

TABLE 8.	BIENTOCANTHUS BREADTH	B-7
TABLE 9.	NASAL BRIDGE BREADTH	B-8
TABLE 10.	BIALAR BREADTH BREADTH	B-9
TABLE 11.	BICHEILION BREADTH	B-10
TABLE 12.	BITRAGION BREADTH	B-11
TABLE 13.	BIZYGOMATIC BREADTH	B-12
TABLE 14.	BIGONIAL BREADTH	B-13
TABLE 15.	SELLION-MENTON LENGTH	B-14
TABLE 16.	SELLION-SUPRAMENTALE LENGTH	B-15
TABLE 17.	SELLION-STOMION LENGTH	
TABLE 18.	SELLION-SUBNASION LENGTH	B-17
TABLE 19.	MIDNASAL BRIDGE HEIGHT FROM MAXILLONASALE PLANE LINE	B-18
TABLE 20.	PRONASALE HEIGHT FROM MAXILLOALARE PLANE LINE	8-19
TABLE 21.	PRONASALE HEIGHT FROM SELLION-PROMENTALE PLANE LINE	B-20
TABLE 22.	SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE	B-21
TABLE 23.	SELLION HEIGHT FROM LATERAL CRBITAL PLANE LINE	B-22

## HEAD AND FACE ANTHROPOMETRY OF ADULT U.S. CIVILIANS

#### INTRODUCTION

This report presents a total of 17 traditional and 5 new head and facial dimensions from a random, composite U.S. female and male civilian population measured over a period of 25 years. The 5 new measurements, identified to describe specific anatomical relationships of the face, can have a direct influence on equipment design concepts. The total survey population includes 367 adult subjects (195 females and 172 males) in the age range of 17 through 69 years representing a predominantly high ratio of Caucasian subjects to subjects of other ethnic origins. The survey population includes subjects used in CAMI experimental projects and selected volunteer subjects; it is hereafter referred to as the CAMI/1992 data base. The data summary format is specifically intended for basic design concept use.

These 22 dimensions describe essential anatomical characteristics for use as basic design criteria in the development of protective equipment for the head and face. Currently, there are standardized, traditional anthropometric data that describe adult two-dimensional face and head characteristics of large military populations. While these data typically describe basic dimensions of major surface features or areas, they do not include specific features and/or combinations of features that are considered essential for design use. The reader should also understand that the population demographics of military data may not necessarily represent comparative civilian populations in age range or distribution of dimensional extremes. The limited availability and use of initial 3-dimensional contour plot data (1) provides excellent design criteria, but is not presently available in a practical or convenient format.

In the practice of descriptive anatomy and anthropometry, multiple labels appear in published format descriptions of identical or associated landmarks and dimensions. Because of the lack of standardization of anatomical and anthropometrical terminology, a user of these types of data and labels can be confused or misled. A thesaurus of selected anatomical and anthropometrical terms, specific to descriptive labels used in this report, is provided as a cross reference for anthropometrical dimensions. This thesaurus is presented in Appendix A.

#### ANTHROPOMETRIC DATA BASES

A review of major anthropometric surveys (ref. 2 through 15) confirms the absence of certain head and face dimensional data for adult U.S. female civilians and limited data for adult U.S. male civilians. Although the most complete military surveys include both female and male data, there are no known civilian population data bases considered adequate for comparative analysis.

The potential overall effect of data bias, if any, in variable population demographics, is not exactly known when using military data to represent civilian values. Typically, physical changes in facial soft tissue can be associated with age and health status. Relative to this type of soft tissue change caused by factors of age and physical conditioning, these changes in anatomical characteristics of the face may be least affected for design needs.

Most current military surveys are also limited in the number and selection of descriptive dimensions of the face. This is a typical problem that is encountered when attempting to make comparisons between most anthropometric data bases. The experimental design for a major population survey usually provides for only those anthropometric dimensions that are specific to the needs of the target population. Other than the traditional measurements of head length and breadth, face breadth, and nose length, there are few consistent dimension selections from one survey to another.

Specific to the need of understanding the relationship of the CAMI/1992 data established in this report with other available data bases, selected key measurements are used for comparisons of 6 face and 2 head dimensions. These key measurements, selected as typical representations of the primary anatomical features, include head

breadth, head length, face breadth, face length, nose breadth, and nose length. These comparisons provide some basic insight to a generalized relationship between the CAMI/1992 data and the selected referenced data (ref. 2 through 14) in gross proportions and size. All reference data, including both international and U.S. population surveys, are considered to be representative of the basic population demographics in size and shape ranges of proportional anatomical variability. These data define the comparative extreme value ranges. Although there are few international female data for specific key comparisons, the international male data values appear to be valid substitutes. It is typical to find small incremental differences in the extreme low value ranges for female and male international data, particularly in the Asiatic-Pacific populations. For the primary intent of this report, the low percentile data of international males will be appropriate for female comparisons. The earliest composite 1946 survey (2, 3) of 8,543 female and 85,000 male army separatees is one of the primary U.S. data resources used to establish the extreme range of dimensional limits for comparison. The 1946 survey may, in fact, closely represent a general civilian population distribution for females and males.

The remaining key reference sources include anthropometric surveys of Iranian (11), Japanese (13), Korean (8), and Vietnamese (5) male military personnel. With the exception of the Iranian survey, the other Asiatic-Pacific surveys of Japanese, Korean, and Vietnamese represent the smallest facial dimensions that are critical to design needs. Comparisons of Asiatic-Pacific populations are considered vital to establish practical extreme range data. Significant changes are occurring in U.S. anthropometry demographics through the increase of Asiatic-Pacific populations. In many instances, current

equipment designs are compromised in functional performance through lack of proper body fit.

Specific key dimensional relationships between the CAMI/1992 and reference data are illustrated by graphs in Figures 1, 2, 3, 4, 5 and 6. These X-Y type graphs plot the data value range as a function of percentile distribution. Selection of reference data for each key dimension (extreme value range) was made on the basis of the lowest and highest value range for females and/or males.

Relative to all key anthropometric dimensions, the CAMI/1992 female data fall within the range limits of data extremes. This type of female distribution is expected, especially with Asiatic-Pacific population data representing some of the smaller face dimension ranges. With the exception of head breadth, the CAMI/1992 data represent the maximum values for data range extremes.

Exceptions to this occur with 95th through 99th percentile values for certain U.S. Army Separatee male data of 1946. The comparative data bases selected for this report represent surveys conducted from 1946 through 1988.

The user of these data should also be aware of the inherent measurement error through observer and instrumentation techniques in soft tissue anthropometry. Valid comparisons of anthropometric surveys should always be made with full awareness of measurement technique and landmark definitions for compatibility. Soft tissue anthropometry is not a precision technique, but should produce acceptable data if surface landmarks are correctly defined and identified.

FIGURE 1. COMPARATIVE HEAD BREADTH PERCENTILES.




FIGURE 2. COMPARATIVE HEAD LENGTH PERCENTILES.




FIGURE 3. COMPARATIVE FACE BREADTH PERCENTILES.





FIGURE 4. COMPARATIVE FACE LENGTH PERCENTILES.



## FIGURE 5. COMPARATIVE NOSE BREADTH PERCENTILES.





FIGURE 6. COMPARATIVE NOSE LENGTH PERCENTILES.



#### STATISTICAL DESCRIPTION

Consistent with the intended purpose of this report, statistical summaries of the described data base include 17 traditional anthropometric dimensions of the head and face that are comparable with measurements of other surveys. The maximum number of data subjects used for any measurement is 195 females and 172 males. The five new measurements are (a) Biorbital Breadth, (b) Sellion-Supramentale Length, (c) Midnasal Bridge Height from a Maxillonasale Plane Line, (d) Pronasale Height from a Sellion-Promentale Plane line, (e) Sellion height from a Lateral Orbital Plane Line.

Each of these measurements is identified and described with the individual data tables by gender. Summary statistics are included with each data table for reference purposes. For consistency with traditional anthropometric data formats intended for use as design reference information, all values are presented in both English and metric value systems.

Descriptive statistics included with each measurement table are the minimum and maximum values, mean, standard error of the mean, standard deviation, coefficient of variation, symmetry, and kurtosis. The standard deviation and variance are calculated as sample or unbiased data to provide a correction factor for sample size.

Because of a wide range in total subjects used for various measurements, a comparative statistical analysis of the data used both the sample (unbiased) and population (biased) methods to calculate standard deviation and variance. Although there appear to be no significant differences between sample and population method calculations, the more conservative sample method may be more realistic for smaller sample sizes.

## DEFINITIONS OF ANATOMICAL LANDMARKS AND PLANES

Definitions of anatomical landmarks and directional reference planes, used to identify surface measurement points, are listed. In most instances, the surface landmark point represents an internal bone feature that is defined as the actual landmark. It is typical of both anatomical and anthropometrical terminology to have duplicate labels for structures and measurements. Refer to the thesaurus listing of anatomical and anthropometrical terminology to confirm certain interchangeable labels.

Paired surface area points define a line in a derived plane from which a perpendicular distance is measured. Surface landmarks and reference planes used for all measurements are indicated in the anatomical illustrations listed in Figures 7 through 13.

ALARE: The most lateral point or area on the lateral surface of the right and left nostrils.

CHEILION: The lateral junction point of the upper and lower lips. The point is correctly positioned with the lips closed and no facial expression. Right and left points.

ECTOCANTHION: The lateral junction point of the upper and lower eyelids of both right and left eyes.

ECTOORBITALE: The most posterior-lateral point on the frontal process of the zygomatic bone. This point approximates the level of defined ECTOCANTHION landmarks.

ENTOCANTHION: The medial junction point of the upper and lower eyelids of both right and left eyes.

GLABELLA: The most anterior midsagittal point on the forehead at the level of the eyebrow ridges.

GONION: the most posterior-inferior midpoint of the GONIAL Angle fo med by the mandibular body and ramus. Right and left points.

LATERAL ORBITAL RIM POINT: A derived point on the anterior-lateral rim of the orbit at a level of the ECTOCANTHUS landmark. The right and left points define the LATERAL ORBITAL PLANE LINE.

LATERAL ORBITAL PLANE LINE: A derived base plane line established for measuring a projection height of SELLION point from the anterior-lateral rim surfaces of the right and left orbits.

MAXILLOALARE: Bilateral points formed by the junction of each maxillary and mid-posterior alar surface areas. These landmarks define the MAXILLOALARE PLANE LINE.

MAXILLOALARE PLANE LINE: A derived base plane line established for measuring a projection height of PRONASALE point away from anterior cheek surfaces.

MAXILLONASALE: Bilateral midpoints along the noseface junction where the anterior maxillary projection starts to form the nose bridge base. These landmarks define the MAXILLONASALE PLANE LINE.

MAXILLONASALE PLANE LINE: A derived base line plane established for measuring a projection height of MIDNASALE point away from anterior cheek surfaces.

MEDIAL CANTHUS PLANE LINE: A derived base plane line established for measuring a projection height of SELLION point. This line plane is coincident with the right and left ENTOCANTHUS landmarks.

MENTON: The most inferior midsagittal point on the chin.

MIDNASALE: A derived midsagittal point on the nasal bridge midway between the SELLION and PRONASALE points.

MIDPUPIL: The center point of both the right and left eye pupils.

MIDSAGITTAL PLANE: A base anatomical plane that divides the head, face, or other bilateral structures into right and left "mirror-image" parts.

PROMENTALE: The most anterior midsagittal point on the chin prominence.

PRONASALE: The most anterior midsagittal point on the nose tip.

SELLION: The point of greatest midsagittal indentation on the nasal bridge immediately inferior to the supraorbital ridge level.

SELLION-PROMENTALE PLANE LINE: A derived base plane line established for measuring a projection height of SELLION point. This plane line is coincident with the right and left ENTOCANTHUS landmarks.

STOMION: A midsagittal point formed at the junction of closed upper and lower lips when relaxed.


SUBNASALE: A midsagittal point at the junction of the external nasal septum and the superior aspect of the upper lip area (Philtrum).

SUPRAMENTALE: The deepest midsagittal point in the horizontal groove between the lower lip and the chin prominence.

TRAGION: A point located at the most anterior aspect of the upper ear notch formed along the superior edge of the Tragus flap. Right and left points.

ZYGION: The most lateral point on the zygomatic arch for both the right and left sides.

## FIGURE 7. FACE & HEAD LANDMARKS.



MENTON

FIGURE 8. FACE & HEAD LANDMARKS.




FIGURE 9. FACE & HEAD MEASUREMENTS.






FIGURE 10. FACE BREADTH MEASUREMENTS.



## FIGURE 11. FACE & HEAD BREADTH MEASUREMENTS. REFERENCE PLANE LINES FOR HEIGHT MEASUREMENTS.



## FIGURE 12. NASAL BRIDGE HEIGHT MEASUREMENTS FROM REFERENCE PLANE LINES.





FIGURE 13. PRONASALE HEIGHT
MEASUREMENTS FROM REFERENCE PLANE
LINES.



### **DEFINITIONS OF ANTHROPOMETRICAL MEASUREMENTS**

By nature, anthropometric instrumentation techniques for measuring deformable soft tissue surfaces do not result in precision measurements. For greatest consistency in reproducing measurement accuracy, soft tissue measurements should be made with only a slight surface contact. Reference numbers are assigned to each anthropometric measurement as indicated in measurement descriptions, illustrations, data summaries, and listed in Table 1.

TABLE 1. REFERENCE NUMBERS OF ANTHROPOMETRIC MEASUREMENTS

MEASUREMENT NUMBER	ANATOMICAL SURFACE REGION AND ANTHROPOMETRIC MEASUREMENT	FIGURE NUMBER
	CRANIUM [ALL]	
1	HEAD CIRCUMFERENCE	NA
2	HEAD BREADTH	11
3	HEAD LENGTH	9
	FACE [BREADTHS]	
4	BIORBITAL BREADTH	10
5	BIECTOCANTHUS BREADTH	10
6	BIPUPIL BREADTH	10
7	BIENTOCANTHUS BREADTH	10
8	NASAL BRIDGE BREADTH	10
9	BIALAR BREADTH	10
10	BICHEILION BREADTH	10
11	BITRAGION BREADTH	11
12	BIZYGOMATIC BREADTH	10
13	BIGONIAL BREADTH	10
	FACE [LENGTHS]	
14	SELLION-MENTON LENGTH	9
15	SELLION-SUPRAMENTALE LENGTH	9
16	SELLION-STOMION LENGTH	9
17	SELLION-SUBNASION LENGTH	9
	NOSE [PROJECTION HEIGHTS]	
18	MIDNASAL BRIDGE HEIGHT FROM MAXILLONASALE PLANE LINE	12
19	PRONASALE HEIGHT FROM MAXILLOALARE PLANE LINE	12
20	PRONASALE HEIGHT FROM SELLION-PROMENTALE PLANE LINE	13
21	SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE	12
22	SELLION HEIGHT FROM LATERAL ORBITAL PLANE LINE	13

HEAD BREADTH: The maximum bilateral distance between right and left sides of the head. (No.2/Fig.11)

HEAD CIRCUMFERENCE: The maximum distance around the head with the tape placed above the eyebrow ridges and positioned over the greatest posterior projection at the back of the head.

HEAD LENGTH: The maximum distance between GLABELLA and the back of the head in the MIDSAG-ITTAL PLANE. (No.3/Fig.9)

BIALAR BREADTH: The bilateral distance between right and left ALARE landmarks of the nose. (No.9/ Fig.10)

BIECTOCANTHUS BREADTH: The bilateral distance between right and left lateral eyelid junctions (ECTOCANTHION landmarks). (No.5/Fig.10)

BIENTOCANTHUS BREADTH: The bilateral distance between right and left medial eyelid junctions (ENTOCANTHION landmarks). (No.7/Fig.10)

BICHEILION BREADTH: The bilateral distance between right and left CHEILION landmarks without facial expression. (No.10/Fig.10)

BIGONIAL BREADTH: The bilateral distance between right and left GONION landmarks at the GONIAL ANGLES of the mandible. (No.13/Fig.10)

BIORBITAL BREADTH: The bilateral distance between right and left lateral orbital surfaces (ECTO-ORBITALE landmarks). (No.4/Fig.10)

BIPUPIL BREADTH: The bilateral distance between right and left pupil centers of the eyes when looking straight ahead. (No.6/Fig.10)

BITRAGION BREADTH: The bilateral distance between right and left TRAGION landmarks of the ears. Measurement obtained with minimum (slight or minimum compression) tissue contact. (No.11/Fig.11)

BIZYGOMATIC BREADTH: The greatest bilateral distance between the most lateral surfaces of the zygomatic arch (ZYGION landmarks). (No.12/Fig.10)

NASAL BRIDGE BREADTH: The bilateral distance between right and left MAXILLONASALE landmarks at the nose bridge base. (No.8/Fig.10)

SELLION-SUBNASALE LENGTH: The midsagittal distance between SELLION and SUBNASALE landmarks of the nose. (No.17/Fig.9)

SELLION-STOMION LENGTH: The midsagittal distance between SELLION and STOMION landmarks with lips closed and teeth in occlusion. (No.16/Fig.9)

SELLION-SUPRAMENTALE LENGTH: The midsagittal distance between SELLION and SUPRA-MENTALE landmarks with the teeth in occlusion. (No.15/Fig.9)

SELLION-MENTON LENGTH: The midsagittal distance between SELLION and MENTON landmarks with the teeth in occlusion. (No.14/Fig.9)

SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE: A perpendicular projection distance of the SELLION landmark away from a line coincident with the bilateral areas just medial to ENTOCANTHION landmarks. (No.21/Fig.12)

SELLION HEIGHT FROM LATERAL ORBITAL PLANE LINE: A perpendicular projection distance of the SELLION landmark away from a line coincident with bota LATERAL ORBITAL RIM POINTS. (No.22/Fig.13)

MIDNASALE HEIGHT FROM MAXILLONASALE PLANE LINE: A perpendicular projection distance of the MIDNASALE landmark away from a line coincident with the right and left MAXILLONASALE landmarks. (No.18/Fig.12)

PRONASALE HEIGHT FROM MAXILLOALARE PLANE LINE: A perpendicular projection distance of the PRONASALE landmark away from a line coincident with right and left MAXILLOALARE LANDMARKS. (No.19/Fig.12)

PRONASALE HEIGHT FROM SELLIONPRO-MENTALE PLANE LINE: The perpendicular projection distance of PRONASALE away from a midsagittal line coincident with the SELLION and PROMENTALE landmarks. (No. 20/Fig. 13)

#### REFERENCES

- Young, J.W., Chandler, R.F., Snow, C.C., et al. Anthropometric and Mass Distribution Characteristics of the Adult Female. Washington, D.C.: 1983; FAA publication no. FAA-AM-83-16, Revised Edition.
- Randall, F.E., Munro, E.H. Reference Anthropometry of Army Women. Report No. 149, Environmental Protection Section, Climatic Research Laboratory, Lawrence, Massachusetts, 1948.
- Newman, R.W., White, R.M. Reference Anthropometry of Army Men. Report No. 180, Environmental Protection Section, Climatic Research Laboratory, Lawrence, Massachusetts, 1951.
- Hertzberg, H.T.E., Daniels, G.S., Churchill, E. Anthropometry of Flying Personnel-1950. Technical Report No. 52-321, Wright Air Development Center, Wright Patterson Air Force Base, Ohio, 1954.
- White, R.M. Anthropometric Survey of the Armed Forces of the Republic of Vietnam. U.S. Army Natick Laboratories, Natick, Massachusetts, 1964.
- Snow, C.C., Snyder, R.G. Anthropometry of Air Traffic Control Trainees. Washington, D.C.: 1965; FAA publication no. AM 65-26.

- Unpublished U.S.A.T. Systems Command Anthropometric Data furnished to Webb Associates, Inc., Yellow Springs, Ohio, by the Aerospace Medical Research Laboratories, Wright Patterson Air Force Base, Ohio, 1965.
- Hart, G.L., Rowland, G.E., Malina, R. Anthropometric Survey of the Armed Forces of the Republic of Korea. Technical Report TR EPT-7, Pioneering Research Laboratory, U.S. Army, Natick Laboratories, Natick, Massachusetts, 1967.
- White, R.M., Churchill, E. The Body Size of Soldiers: U.S. Army Anthropometry-1966. Technical Report No. 72-51-CE, U.S. Army Natick Laboratories, Natick, Massachusetts, 1971.
- 10. Unpublished U.S.A.F. Systems Command Anthropometric Data of flying personnel furnished to Webb Associates, Inc., Yellow Springs, Ohio, by the Aerospace Medical Research Laboratories, Wright Patterson Air Force Base, Ohio, 1967.
- Neorani, S.E., Dillars, C.N. Jr. Anthropometric Survey of the Imperial Iranian Armed Forces, Data Collection and Analysis, Volumes I and II. Technical Report of the Combat Research and Evaluation Center, Imperial Iranian Ground Forces, Teheran, Iran, 1970.
- Clauser, C.E. et al. Anthropometry of Air Force Women. Technical Report No. 70-5, Aerospace Medical Laboratories, Wright Patterson Air Force Base, Ohio, 1972.
- 13. Yokohori, E. Anthropometry of JASDF Personnel and Its Application for Human Engineering. Aeromedical Laboratory, Japanese Air Self Defense Force, Tachikawa Air Base, Toyko, Japan, 1972.
- Donelsor, S.M., Gordon, C.C. 1988 Anthropometric Survey of U.S. Army Personnel: Pilot Summary Statistics. Technical Report Natick/TR-89/044. U.S. Army Natick RD&E Center, Natick, Massachusetts. 1991.

### APPENDIX A

## **ANTHROPOMETRY THE SAURUS**

This appendix is a thesaurus of selected anthropometrical measurement terms that use multiple labels to identify specific dimensions of superficial head and face structures. The terms listed as REPORT TERMS are those used in this report. Equivalent terms, typical of those found in many published reports, are listed as ALTERNATE TERMS:

REPORT TERMS BIALAR BREADTH	ALTERNATE TERMS BIALAR DIAMETER  NASAL BREADTH
	NOSE BREADTH
BICHELION BREADTH	
	MOUTH BREADTH
	LIP LENGTH
BIECTOCANTHUS BREADTH	RIOCIII AD RDEADTH
DIECTOCANTITOS BREADTIT	BIOCULAR BREADTH
	MAXIMUM BIOCULAR DIAMETER
	LATERAL CANTHUS DIAMETER
	Bilbidib Callitoo Diawibi bic
BIENTOCANTHUS BREADTH	INTEROCULAR BREADTH
	INTEROCULAR DIAMETER
	MEDIAL CANTHUS DIAMETER
BIGONIAL BREADTH	DICONIAL DIAMETER
DIGUNIAL DREADITI	MAXIMUM LOWER FACE BREADTH
	MARINON LOWER THOE BREADTH
BIORBITAL BREADTH	-MAXIMUM FRONTAL DIAMETER
	MAXIMUM ORBITAL DIAMETER
	MAXIMUM BIORBITAL BREADTH
BIPUPIL BREADTH	INTERPLIPILLARY BREADTH
DIL OTTE BROWN TITE	INTERPUPILLARY DIAMETER
BIZYGOMATIC BREADTH	FACE BREADTH
	UPPER FACE BREADTH
HEAD BREADTH	
	MAXIMAL HEAD BREADTH
	MAXIMAL HEAD DIAMETER
HEAD LENGTH	HEAD LENGTH
	GLABELLAR-OPISTHOCRANION LENGTH

REPORT TERMS ALTERNATE TERMS MIDNASAL BRIDGE HEIGHT FROM MAXILLONASALE PLANE LINE ----------NASAL BRIDGE HEIGHT NASAL BRIDGE SALIENT NASAL BRIDGE BREADTH -----NOSE BRIDGE BREADTH PRONASALE HEIGHT FROM MAXILLONASALE PLANE LINE -----NOSE PROJECTION HORIZONTAL NOSE PROTRUSION NASAL TIP SALIENT SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE ------- NASAL ROOT SALIENT NASAL ROOT HEIGHT SELLION-MENTON LENGTH ------ MENTON-SELLION LENGTH FACE LENGTH TOTAL FACE LENGTH **FACE HEIGHT** SELLION-SUBNASION LENGTH-SUBNASALE-SELLION LENGTH SUBNASALE-NASAL ROOT DEPRESSION LENGTH NASAL HEIGHT **NOSE HEIGHT NOSE LENGTH** 

### APPENDIX B

MALE

TABLE 2. HEAD CIRCUMFERENCE

FEMALE

	<del></del>		
SUMMARY :	STATISTICS	SUMMARY STATIS	TICS
MILLIMETERS	INCHES	MILLIMETERS	INCHES

549.84 1.88 20.48	MEAN STD ERROR(MEAN) STD DEVIATION	21.65 Ø.Ø7 3.71	574.54 MEAN 1.55 STD ERROR(MEAN) 16.22 STD DEVIATION	22.62 Ø.Ø6 Ø.64
5Ø6.98 614.93	MINIMUM MAXIMUM	19.96 24.21	541.02 MINIMUM 620.01 MAXIMUM	21.30
	OF VARIATION(%)	3.71 Ø.54 3.37	COEFF. OF VARIATION(%) SYMMETRY KURTOSIS	2.81 Ø.24 2.39
NUMBER (	OF SUBJECTS	119	NUMBER OF SUBJECTS	11Ø
	PERCENTILES		PERCENTILES	
MILLIMET	ers	INCHES	MILLIMETERS	INCHES
5Ø6.98	1 ST	19.96	547.88 1 ST	21.57

MILLIMETERS			INCHES	MILLIMETERS			INCHES
5Ø6.98	1	ST	19.96	547.88	1	ST	21.57
518.92	5	$\mathbf{TH}$	20.43	549.91	5	TH	21.65
525.Ø2	1Ø	TH	2Ø.67	554.99	10	TH	21.85
534.92	25	TH	21.Ø6	56Ø.Ø7	25	TH	22.05
547.88	5Ø	TH	21.57	573.Ø2	5Ø	TH	22.56
561.Ø9	75	TH	22.Ø9	586.99	75	TH	23.11
574.Ø4	9Ø	TH	22.6Ø	595.12	9Ø	TH	23.43
581.91	95	TH	22.91	599.95	95	TH	23.62
6Ø5.Ø3	99	TH	23.82	6Ø8.Ø8	99	ΤH	23.94

HEAD CIRCUMFERENCE: The maximum distance around the head with the tape placed above the eyebrow ridges and positioned over the greatest posterior projection at the back of the head.

TABLE 3. HEAD BREADTH

MALE

ST	MARY STATISTICS	}	S	MARY STATISTIC	5
MILLIME	TERS	INCHES	MILLIME	TERS	INCHES
145.76	MEAN	5.74	152.39	MEAN	6.00
Ø.43	STD ERROR (MEAN)	Ø.Ø2	Ø.41	STD ERROR (MEAN	0.02
5.57	SID DEVIATION	Ø.22	5.34	STD DEVIATION	
128.Ø2	MINIMUM	5.Ø4	138.94	MINIMIM	5.47
161.Ø4	MAXIMUM	6.34	166.12	MAXIMIM	6.54
COEFF.	OF VARIATION(%)	3.8Ø	COEFF.	OF VARIATION(%)	3.5Ø
SYMMETE	RY	Ø.Ø2	SYMET	RY	-0.01
KURTOSI	ts	3.25	KURTOS]	<u>s</u>	2.48
NUMBER	OF SUBJECTS	169	NUMBER	OF SUBJECTS	167
	PERCENTILES			PERCENTILES	
MILLIMET	TERS	INCHES	MILLIMET	TERS	INCHES
131.Ø6	1 ST	5.16	140.97	1 ST	5.55
136.91	5 TH	5.39	144.02	5 TH	5.67
137.92	10 TH	5.43	145.Ø3	10 TH	5.71
141.99	25 TH	5.59	148.Ø8	25 TH	5.83
146.Ø5	5Ø TH	5.75	152.91	5Ø TH	6.Ø2
149.10	75 TH	5.87	155.96	75 TH	6.14
152.91		6.Ø2	159.00	9Ø TH	6.26
154.94	95 TH	6.10	161.04	95 TH	6.34
157.99	99 TH	6.22	163.07	99 TH	6.42
152.91	9Ø TH	6.Ø2 6.1Ø	159.00 161.04	9Ø TH 95 TH	6.2 6.3

HEAD BREADTH: The maximum bilateral distance between the right and left sides of the head.

TABLE 4. HEAD LENGTH

MALE

SUM	MARY STATISTICS	5		SU	mmary statisti	cs
MILLIMETE	RS	INCHES		MILLIMET	ERS	INCHES
187.84	MEAN	7.40		199.57	MEAN	7.86
Ø.56	SID ERROR (MEAN)	Ø.Ø2		ø.59	STD ERROR (MEA)	N) Ø.Ø2
7.29	STD DEVIATION	Ø.29		7.56	STD DEVIATION	N Ø.3Ø
165.10	MINIMUM	6.5Ø		180.09	MINIMUM	7.Ø9
216.92	MAXIMUM	8.54		223.Ø1	MAXIMUM	8.78
COEFF. O	F VARIATION(%)	3.86		COEFF.	OF VARIATION(%	) 3.76
SYMMETRY		Ø.21		SYMMETR	Y	Ø.32
KURTOSIS		4.32		KURTOST	S	3.62
NUMBER O	F SUBJECTS	169		NUMBER	OF SUBJECTS	166
	PERCENTILES				PERCENTILES	
MILLIMETE	RS	INCHES	***************************************	MILLIMET	ERS	INCHES
17Ø.94	1 ST	6.73		182.12	1 ST	7.17
175.Ø1	5 TH	6.89		187.96	5 TH	7.4Ø
179.Ø7	1Ø TH	7.Ø5		191.Ø1	1Ø TH	7.52
182.88	25 TH	7.2Ø		194 <i>-</i> Ø6	25 TH	7.64
187.96	5Ø TH	7.4Ø		198.88	5Ø TH	7.83
193.Ø4	75 TH	7.6Ø		2Ø3.96	75 TH	8.Ø3
196.Ø9	9Ø TH	7.72		2Ø9.Ø4	9Ø TH	8.23
198 - 12	95 TH	7.8Ø		213.11	95 TH	8.39
2Ø3.96	99 TH	8.Ø3		217.93	99 TH	8.58

HEAD LENGTH: The maximum distance between GLABFILLA and the back of the head in the MIDSAGITTAL PLANE.

TABLE 5. BIORBITAL BREADTH

<del></del>					
SUM	MARY STATISTICS	}	su	Mary Slatistic	3
MILLIMETE	RS	INCHES	MILLIMET	es Es	INCHES
113.79	MEAN	4.48	119.01	MEAN	4.69
Ø.43	SID ERROR (MEAN)	Ø. <b>Ø</b> 2	Ø.5Ø	STD ERROR (MEAN)	9.02
5.Ø\$	SID DEVIATION	Ø.2Ø	5.81	STD DEVIATION	Ø. 23
102.11	MINIMLM	4.02	99.06	MINIMM	3.9Ø
128.Ø2	MAXIMUM	5.Ø4	134.87	MAXIMIM	5.32
COEFF. O	F VARIATION(%)	4.42	COEFF.	OF VARIATION(%)	4.88
SYMMETRY	•	Ø.31	SYMMETR	Y	-Ø.39
KURTOSIS		2.76	KURTOSI	S	3.98
NUMBER O	F SUBJECTS	140	NLM3EK	OF SUBJECTS	137
	PERCENTILES			PERCENTILES	
MILLIMETE	RS	INCHES	MILLIMET	ERS	INCHES
102 11	1 ST	4.Ø2	99.Ø6	1 ST	3.9Ø
1Ø5.92	5 TH	4.17	107.95	5 TH	4.25
1Ø7.95	1Ø TH	4.25	112.Ø1	10 TH	4.41
1Ø9.98	25 TH	4.33	115.06	25 TH	4.53
113.Ø3	5Ø TH	4.45	119.13	50 TH	4.69
118.11	75 TH	4.65	122.94	75 TH	4.84
119.89	9Ø TH	4.72	125.98		4.96
121.92	95 TH	4.8Ø	128.Ø2	95 TH	5.Ø4
125.98	99 TH	4.96	133.1Ø	99 TH	5.24

BIORBITAL EREADTH: The bilateral distance between the right and left lateral orbital surfaces (ECTCORBITALE landmarks).

TABLE 6. BIECTOCANTHUS EREADTH

MALE

	· · · · · · · · · · · · · · · · · · ·					
ST	MMARY STATISTICS	5	St	immary st	PATISTIC	5
CENTIME	TERS	INCHES	CENTIMET	TERS		INCHES
88.53	MEAN	3.49	91.31	М	EAN	3.6Ø
Ø.41	STD ERROR (MEAN)	Ø.Ø2	Ø.5Ø	STD ER	ROR (MEAN)	Ø.Ø2
4.78	STD DEVIATION	Ø. 19	4.99	STD DI	EVIATION	Ø.2Ø
76.96	MINIMUM	3.Ø3	82.Ø4	MIN	MIMIN	3.23
1Ø3.89	MAXIMUM	4.Ø9	107.95	MAX	CIMUM	4.25
COEFF.	OF VARIATION(%)	5.41	COEFF.	OF VARIA	TION(%)	5.43
SYMMETE	ΥY	Ø. 44	SYMMETE	Ϋ́		Ø.57
KURTOSI	rs .	3.Ø1	KURTOSI	S		3.41
NUMBER	OF SUBJECTS	134	NUMBER	OF SUBJ	ECTS	1Ø1
	PERCENTILES			PERCEI	VTILES	
CENTIMET	TERS	INCHES	CENTIMET	ERS		INCHES
76.96	1 ST	3.Ø3	82.Ø4	1	ST	3.23
82.Ø4	5 TH	3.23	84.Ø7	5	TH	3.31
83.Ø6	10 TH	3.27	84.58	1Ø	TH	3.33
85.Ø9	25 TH	3.35	87.88	25	TH	3.46
87.88	5Ø TH	3.46	9Ø.93	5Ø	TH	3.58
9Ø. 93	75 TH	3.58	93.98	75	TH	3.7Ø
95. <i>0</i> Ø	9Ø TH	3.74	98.Ø4	9Ø	TH	3.86
97.Ø3	95 TH	3.82	1Ø1.Ø9		TH	3.98
100.08	99 TH	3.94	1Ø3.12	99	TH	4.06

BIECTOCANTHUS BREADTH: The bilateral distance between the right and left lateral eyelid junctions (ECTOCANTHION landmarks).

TABLE 7. BIPUPIL BREADTH

st	MARY STATISTICS	3	SU	MMARY STATISTIC	s
MILLIME	TERS	INCHES	MILLIMET	<b>er</b> s	INCHES
58.44	MEAN	2.30	61.39	MEAN	2.42
Ø.28	SID ERROR (MENN)	Ø.Ø1	Ø.36	STD ERROR (MEAN	r) Ø.Ø1
3.32	STD DEVIATION	Ø. 13	<b>3.6</b> 3	STD DEVIATION	Ø. 14
52.Ø7	MINIMUM	2.Ø5	54.99	MINIMUM	2.17
68.Ø7	MAXIMIM	2.68	7Ø.99	MAXIMUM	2.80
COEFF.	OF VARIATION(%)	5.66	COEFF.	OF VARIATION(%)	5.88
SYMMET	RY ·	Ø.5Ø	SYMMETE	Y	Ø.55
KURTOS I	s	2.85	KURTOSI	S	2.99
NUMBER	OF SUBJECTS	136	NUMBER	of subjects	1Ø2
	PERCENTILES			PERCENTILES	
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
52.Ø7	1 ST	2.05	54.99	1 <b>ST</b>	2.17
54.10	5 TH	2.13	56.01	5 TH	2.20
54.61	10 TH	2.15	57.00	10 TH	2.24
55.88	25 TH	2.205	59.00	25 TH	2.32
57.91	5Ø TH	2.28	61.21	5Ø TH	2.40
59.94	75 TH	2.36	62.99	75 TH	2.48
62.99	9Ø TH	2.48	65.99	9Ø TH	2.60
64.Ø1	95 TH	2.52	68.00	95 TH	2.68
68.Ø7	99 TH	2.68	7Ø.99	99 TH	2.80

BIPUPIL BREADIH: The bilateral distance between the right and left pupil centers of the eyes when looking straight ahead.

TABLE 8. BIENTOCANTHUS BREADTH

MALE

St	MMARY STATISTICS		. Su	MMARY STATISTIC	cs .
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
3ø.57		1.2Ø	32.Ø8	MEAN	1.26
Ø.23	,	·- • •	Ø.27	SID ERROR (MEAN	N) Ø.Ø1
2.71	SID DEVIATION	Ø. 11	2.95	STD DEVIATION	Ø. 12
23.88	MINIMUM	Ø. 95	25.91	MINIMUM	1.Ø2
39.89	MAXIMUM	1.58	39.88	MAXIMUM	1.58
COEFF.	OF VARIATION(%)	8.72	COEFF.	OF VARIATION(%)	9.ø3
SYMMETE	RY	Ø.25	SYMMETR	Y	Ø.52
KURTOSI	rs .	4.04	KURTOSI	S	2.76
NUMBER	OF SUBJECTS	145	NUMBER	OF SUBJECTS	115
	PERCENTILES			PERCENTILES	
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES
23.88	1 ST	Ø. 94	25.91	1 ST	1.Ø2
25.91	5 TH	1.Ø2	27.94	5 TH	1.10
26.92	1Ø TH	1.Ø6	28.96	1Ø TH	1.14
28.96	25 TH	1.14	29.97	25 TH	1.18
29.97	5Ø TH	1.18	32.00	5Ø TH	1.26
32.00	75 TH	1.26	34.04	75 TH	1.34
34.04	9Ø TH	1.34	36.Ø7	90 TH	1.42
35.Ø5	95 TH	1.38	37.Ø8	95 TH	1.46
39.12	99 TH	1.54	39.12	99 TH	1.54

BIENTOCANTHUS BREADTH: The bilateral distance between the right and left medial eyelid junctions (ENTOCANTHION landmarks).

TABLE 9. NASAL BRIDGE BREADTH

FEMALE	MALE

			· · · · · · · · · · · · · · · · · · ·	<del></del>	
SU	mmary statistic	s	su	MMARY STATISTIC	S
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES
31.83	MEAN	1.25	33.17	MEAN	1.31
Ø.21	STD ERROR (MEAN	) Ø.Ø1	Ø.25	STD ERROR (MEAN	
2.72	SID DEVIATION	0.11	2.99	STD DEVIATION	
23.11	MINIMIM	Ø. 91	25.91	MINIMIM	1.Ø2
39.88	MAXIMUM	1.58	39.12	MIMIXAM	1.54
COEFF.	OF VARIATION(%)	8.42	COEFF.	OF VARIATION(%)	8.86
SYMMETE	RY .	Ø.Ø3	SYMMETR	Y	-Ø.34
KURTOSI	S	3.45	KURTOSI	S	2.62
NUMBER	OF SUBJECTS	174	NUMBER	OF SUBJECTS	145
	PERCENTILES			PERCENTILES	
MILLIMET	ERS	INCHES	MILLIMET	ERS	INCHES
24.89	1 ST	Ø. 98	25.91	1 ST	1.Ø2
26.92	5 TH	1.06	27.94	5 TH	1.10
28.96	10 TH	1.14	28.96	10 TH	1.14
29.97	25 TH	1.18	3Ø.99	25 TH	1.22
32.ØØ	5Ø TH	1.26	33.Ø2	5Ø TH	1.30
33.Ø2	75 TH	1.30	35.Ø5	75 TH	1.38
35.Ø5	9Ø TH	1.38	37.Ø8	90 TH	1.46
36.Ø7	95 TH	1.42	38.1Ø	95 TH	1.50
38. 1Ø	99 TH	1. <i>50</i>	39.12	99 TH	1.54

NASAL BRIDGE BREADTH: The bilateral distance between the right and left MAXILLONASALE landmarks at the nose bridge base.

TABLE 10. BIALAR BREADTH

MALE

	<del></del>				<del></del>
SUMMARY STATISTICS			SUMMARY STATISTICS		
MILLIMET	ERS	INCHES	MILLIMET	ERS	INCHES
33.29	MEAN	1.31	36.78	MEAN	1.45
Ø.24	SID ERROR (MEAN	Ø.Ø1	Ø.28	STD ERROR (MEA	N) Ø.01
3.16	STD DEVIATION	Ø. 13	3.42	SID DEVIATIO	N Ø. 14
27.00	MINIMIM	1.Ø6	3Ø <b>.</b> 99	MINIMUM	1.22
43.00	MAXIMIM	1.69	50.04	MAXIMIM	1.97
COEFF.	OF VARIATION(%)	9.56	COEFF.	OF VARIATION(%	) 9.33
SYMMETR	Y	1.02	SYMMETR	Y	Ø. 92
KURTOSI	S	4.44	KURTOSI	S	4.37
NUMBER	OF SUBJECTS	172	NUMBER	OF SUBJECTS	145
	PERCENTILES			PERCENTILES	
MILLIMET	ERS	INCHES	MILLIMET	ers	INCHES
27.94	1 ST	1.10	3ø. 99	1 ST	1.22
28.96	5 TH	1.14	32.00	5 TH	1.26
29.97	1Ø TH	1.18	32.51	10 TH	1.28
32.00	25 TH	1.26	35.Ø5	25 TH	1.38
33.Ø2	5Ø TH	1.3Ø	36.Ø7	5Ø TH	1.42
35.Ø5	75 TH	1.38	38.10	75 TH	1.50
37.Ø8	9Ø TH	1.46	40.89	9Ø TH	1.61
39,88	95 TH	1.57	42.93	95 TH	1.69
42.93	99 TH	1.69	48.Ø1	99 TH	1.89

BIALAR BREADTH: The bilateral distance between the right and left ALARE landmarks of the nose.

TABLE 11. BICHEILION BREADTH

***		· · · · · · · · · · · · · · · · · · ·			
SUM	MARY STATISTICS		st	MMARY STATISTIC	s
MILLIMETE	RS	INCHES	MILLIME	TERS	INCHES
48.25	MEAN	1.90	52.23	MEAN	2.Ø6
Ø.33	SID ERROR (MEAN)	Ø.Ø1	Ø.36	STD ERROR (MEAN	0.01
3.95	SID DEVIATION	Ø.21	3.91	STD DEVIATION	
37.Ø8	MINIMUM	1.46	42.93	MINIMIM	1.69
56.9Ø	MAXIMUM	2.24	62.99	MAXIMUM	2.48
COEFF. O	F VARIATION(%)	8.Ø6	COEFF.	OF VARIATION(%)	7.47
SYMMETRY		ø.ø5	SYMMET	ZY.	-0.01
KURTOSIS		2.65	KURTOS1		3.13
NUMBER O	F SUBJECTS	145	NUMBER	OF SUBJECTS	115
	PERCENTILES	•		PERCENTILES	
MILLIMETE	RS	INCHES	MILLIMET	TERS	INCHES
37.Ø8	i st	1.46	42.93	1 ST	1.69
41.91	5 TH	1.65	43.94	5 TH	1.73
42.93	10 TH	1.69	48.Ø1	10 TH	1.89
44.96	25 TH	1.77	50.04	25 TH	1.97
48.Ø1	5Ø TH	1.89	52.07	5Ø TH	2.Ø5
51.Ø5	75 TH	2.Ø1	55.12	75 TH	2.17
53.Ø9	9Ø TH	2.Ø9	56.90		2.24
55.12	95 TH	2.17	57.91	95 TH	2.28
56.90	99 TH	2.24	6Ø.96	99 TH	2.40

BICHEILION BREADTH: The bilateral distance between the right and left CHEILION landmarks without facial expression.

TABLE 12. BITRAGION BREADTH

MALE

su	MMARY STATISTICS		SU	MMARY STATISTICS	5
MILLIMET	ers	INCHES	MILLIMET	ERS	INCHES
131.70	MEAN	5.19	14Ø.89	MEAN	5.55
Ø.5Ø	STD ERROR (MEAN)	Ø.Ø2	ø.58	STD ERROR (MEAN)	Ø.Ø2
5.47	STD DEVIATION	Ø.22	5.75	SID DEVIATION	Ø.23
119.13	MINIMIM	4.69	128.Ø2:	MINIMIM	5.Ø4
144.Ø2	MAXIMUM	5.67	152.91	MAXIMIM	6.Ø2
COEFF.	OF VARIATION(%)	4.14	COEFF.	OF VARIATION(%)	4.Ø5
SYMMETR	Y	Ø. 14	SYMMETR		-0.04
KURTOSI	S	2.49	KURTOSI	s. 🗇	2.25
NUMBER (	OF SUBJECTS	12Ø	NUMBER	OF SUBJECTS	98
	PERCENTILES			PERCENTILES	
MILLIMET	ers	INCHES	MILLIMET	ERS	INCHES
119.13	1 ST	4.69	128.Ø2	1 ST	5.04
121.92	5 TH	4.80	132.08	5 TH	5.20
123.95	10 TH	4.88	133.10	1Ø TH	5.24
127.00	25 TH	5.ØØ	136.40	25 TH	5.37
132.Ø8	5Ø TH	5.2Ø	141.99	50 TH	5.59
134.87	75 TH	5.31	145.03	75 TH	5.71
138.94	9Ø TH	5.47	149.10	9Ø TH	5.87
140.97	95 TH	5.55	150.11	95 TH	5.91
143.ØØ	99 TH	5.63	151.91	99 TH	5.98

BITRAGION BREADIH: The bilateral distance between the right and left TRAGION landmarks of the ears. Measurement obtained with minimum (no compression) tissue contact.

TABLE 13. BIZYGOMATIC BREADTH

MALE

······································		<del></del>	<del></del>			
SU	SUMMARY STATISTICS			SUMMARY STATISTICS		
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES	
131.81	MEAN	5.18	140.67	MEAN	5.54	
Ø.36	STD ERROR (MEAN	) Ø.Ø1	Ø.46	STD EPROR (MEAN)	Ø.Ø2	
5.Ø9	STD DEVIATION	Ø.2Ø	6.02	SID DEVIATION		
119.12	MINIMIM	4.68	125.98	MINIMUM	4.96	
148.Ø8	MAXIMIM	5.83	152.91	MAXIMUM	6.02	
COEFF.	OF VARIATION(%)	3.85	COEFF.	OF VARIATION(%)	4.26	
SYMMETR	Y.	Ø.12	SYMMETR	Y	-Ø.Ø8	
KURTOSI	:S	3.11	KURTOSI	S	2.53	
NUMBER	OF SUBJECTS	195	NUMBER	OF SUBJECTS	171	
	PERCENTILES			PERCENTILES		
MILLIMET	ERS	INCHES	MILLIMET	ERS	INCHES	
119.89	1 ST	4.72	127.00	1 S2	5.00	
122.94	5 TH	4.84	130.05	5 TH	5.12	
124.97	10 TH	4.92	133.1Ø	10 TH	5.24	
128.Ø2	25 TH	5.94	135.89	25 TH	5.35	
132.Ø8	50 TH	5.20	139.95	5Ø TH	5.51	
134.87	75 TH	5.31	145.Ø3	· 75 TH	5.71	
137.92	9Ø TH	5.43	149.10	9Ø TH	5.87	
139.95	95 TH	5.51	150.11	95 TH	5.91	
145.Ø3	99 TH	5.71	151.89	99 TH	5.98	
				· ·		

BIZYGOMATIC BREADTH: The greatest bilateral distance between the lateral cheek surfaces of the zygomatic arch (ZYGION landmarks).

TABLE 14. BIGONIAL BREADTH

		<u></u>			
SU	MMARY STATISTICS	5	SU	MARY STATISTIC	5
MILLIMET	ERS	INCHES	MILLIMET	ERS	INCHES
100.87	MEAN	3.97	108.50	MEAN	4.27
Ø.41	STD ERROR (MEAN)	Ø.Ø2	Ø.59	STD ERROR (MEAN)	0.02
5.40	STD DEVIATION	Ø.21	7.05	STD DEVIATION	Ø.28
84.Ø7	MINIMUM	3.31	د ۶.۰۶	MINIMIM	3.58
117.Ø9	MAXIMIM	4.61	133.10		5.24
COEFF.	OF VARIATION(%)	5.34	COEFF.	OF VARIATION(%)	6.47
SYMMETR	Y	Ø. 15	SYMMETR	Y	Ø.43
KURTOSI	S	3.22	KURTOSI	S	3.54
NUMBER	OF SUBJECTS	174	NUMBER (	OF SUBJECTS	144
	PERCENTILES			PERCENTILES	
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES
87.12	1 ST	3.43	90.93	1 ST	3.58
91.95	5 TH	3.62	97.Ø3	5 TH	3.82
92.96	10 TH	3.66	100.08	10 TH	3.94
97.Ø3	25 TH	3.82	103.89	25 TH	4.09
101.09	50 TH	3.98	1Ø7.95	50 TH	4.25
1Ø3.89	75 TH	4.Ø9	113.03	75 TH	4.45
107.95	9Ø TH	4.25	117.09		4.61
109.98	95 TH	4.33	119.13		4.69
112.Ø1	99 TH	4.41	128.02	99 TH	5.04

BIGONIAL EREADIH: The bilateral distance between the right and left GONION landmarks at the GONIAL ANGLES of the mandible.

TABLE 15: SELLION-MENTON LENGTH

FEMALE	MALE
--------	------

. <del></del>					
st	MMARY STATISTICS	5	SU	MMARY STATISTIC	×
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
110.97	MEAN	4.37	12Ø.91	MEAN	4.76
Ø.46	STD ERROR (MEAN)	) Ø.Ø2	Ø.56	STD ERROR (MEAN	i) Ø.56
6.47	STD DEVIATION	Ø.25	7.36	SID DEVIATION	Ø.29
93.98	MINIMIM	3.70	1Ø5.92	MINIMIM	4.17
133.10	MAXIMUM	5.24	139.95	MAXIMUM	5.51
COEFF.	OF VARIATION(%)	5.8Ø	COEFF.	OF VARIATION(%)	6.07
SYMMETE	XX	Ø.27	SYMMETR	Y	Ø.Ø9
KURIOSI	is .	3.33	KURTOSI	S	2.57
NUMBER	OF SUBJECTS	195	NUMBER	OF SUBJECTS	172
	PERCENTILES		٠	PERCENTILES	
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
95.00	1 ST	3.74	1Ø5.92	1 ST	4.17
100.08	5 TH	3.94	108.97	5 TH	4.29
103.12	1Ø TH	4.06	1Ø9.98	10 TH	4.33
105.92	25 TH	4.17	116.Ø8		4.57
111.00	50 TH	4.37	120.90	5Ø TH	4.76
115.06	75 TH	4.53	125.98		4.96
119.13	9Ø TH	4.69	130.05	9Ø TH	5.12
120.90	95 TH	4.76	132.Ø8	95 TH	5.20
122.94	99 TH	4.84	136.91	99 TH	5.39

SELLION-MENTON LENGTH: The midsagittal distance between the SELLION and MENTON landmarks with the teeth in occlusion.

TABLE 16. SELLION-SUPRAMENTALE LENGTH

·		·········			<del> </del>
St	MMARY STATISTICS	<b>,</b>	su	MMARY STATISTICS	5
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
85.68	MEAN	3.37	93.2Ø	MEAN	3.67
Ø.48	SID ERROR (MEAN)	Ø.Ø2	Ø.62	SID ERROR (MEAN)	Ø.Ø2
5.78	STD DEVIATION	Ø.23	6.66	SID DEVIATION	Ø.26
71.88	MINIMIM	2.84	8ø.ø1	MINIMUM	3.15
107.95	MAXIMM	4.25	1Ø9.98		4.33
COEFF.	OF VARIATION(%)	6.73	COEFF.	OF VARIATION(%)	7.12
SYMMEXE	RY.	Ø.41	SYMMETR	Y	Ø. 17
KURTOS I	S	3.69	KURTOSI	S	2.51
NUMBEP.	OF SUBJECTS	145	NUMBER	OF SUBJECTS	115
	PERCENTILES			PERCENTILES	
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
71.88	1 ST	2.83	8Ø.Ø1	1 ST	3.15
75.95	5 TH	2.99	83.Ø6	5 TH	3.27
78.99	10 TH	3.11	84.Ø7	10 TH	3.31
82.Ø4	25 TH	3.23	88.9Ø	25 TH	3.5Ø
85.Ø9	50 TH	3.35	92.96	5Ø TH	3.66
88.90	75 TH	3.5Ø	97.Ø3	75 TH	3.82
92.96	SØ TH	3.66	101.09		3.98
95.00	95 TH	3.74	104.90		4.13
99.Ø6	99 TH	3.9Ø	1Ø7.95	99 TH	4.25

SELLION-SUPRAMENTALE LENGTH: The midsagittal distance between the SELLION and SUPRAMENTALE landmarks with the teeth in normal occlusion.

TABLE 17. SELLION-STOMION LENGTH

TOTAL AT TO	3/27 77
FEMALE	MALE

	<del></del>	<del></del>		<del>,, , , , , , , , , , , , , , , , , , ,</del>			
SUMMARY STATISTICS		5	st	mary statistic	(CS		
MILLIME	TERS	INCHES	MILLIMET	ers	INCHES		
69.Ø1		2.72	75.7Ø				
	STD ERROR (MEAN)		Ø.44				
4.40	STD DEVIATION	Ø. 18	4.71	STD DEVIATION	N Ø.19		
55.88	MINIMM	2.21	65.Ø2	MINIMUM	2.56		
83.Ø6	MAXIMUM	3.27	90.93	MAXIMUM	3.58		
COEFF.	OF VARIATION(%)	6.45	COEFF.	OF VARIATION(%	6.20		
SYMMETE	RY	Ø. 16	SYMMETE	Y	Ø.25		
KURTOS1	rs .	3.11	KURTOSI	S	3.15		
NUMBER	OF SUBJECTS	145	NUMBER	OF SUBJECTS	115		
	PERCENTILES			PERCENTILES			
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES		
55.88	1 ST	2.20	65.Ø2	1 ST	2.56		
61.98	5 TH	2.44	68.Ø7	5 TH	2.68		
62.99	10 TH	2.48	69.Ø9	1Ø TH	2.72		
66.Ø4	25 TH	2.6Ø	71.88	25 TH	2.83		
69.Ø9	5Ø TH	2.72	74.93	5Ø TH	2.95		
71.88	75 TH	2.83	77.98	75 TH	3.07		
74.93	9Ø TH	2.95	81.Ø3	9Ø TH	3.19		
75.95	95 TH	2.9 <del>9</del>	83.Ø6	95 TH	3.27		
8Ø.Ø1	99 TH	3.15	86.11	99 TH	3.39		

SELLION-STOMION LENGTH: The midsagittal distance between the SELLION and STOMION landmarks with the lips closed and teeth in occlusion.

TABLE 18. SELLION-SUBNASION LENGTH

			<u> </u>		
St	MMARY STATISTICS	5	S	UMMARY STATISTIC	S
MILLIMET	TERS	INCHES	MILLIME	TERS	INCHES
5ø. 18	MEAN	1.93	54.Ø1	MEAN	2.13
Ø.3Ø	SID ERROR (MEAN)	Ø.Ø1	Ø.34	STD ERROR (MEAN	) Ø.34
3.83	STD DEVIATION	Ø. 15	4.09	SID DEVIATION	Ø. 16
38.10	MINIMIM	1.50	41.91	MENEMEM	1.65
62.99	MAXIMIM	2.48	62.89	MUMIXAM	2.48
COEFF.	OF VARIATION(%)	7:56	COEFF.	OF VARIATION(%)	7.60
SYMMETR	Y	Ø.Ø1	SYMMET		-Ø.Ø8
KURTOSI	S	3.31	KURTOS	<u>rs</u>	2.67
NUMBER	OF SUBJECTS	175	NUMBER	OF SUBJECTS	145
	PERCENTILES			PERCENTILES	
MILLIMET	ERS	INCHES	MILLIME	TERS	INCHES
40.89	1 ST	1.61	41.91	1 ST	1.65
43.94	5 TH	1.73	46.99	5 TH	1.85
44.96	10 TH	1.77	49.Ø2	10 TH	1.93
46.99	25 TH	1.85	51.Ø5	25 TH	2.01
50.04	5Ø TH	1.97	- 54 . 1Ø	5Ø TH	2.13
53.Ø9	75 TH	2.Ø9	56.90	75 TH	2.24
55.12	9Ø TH	2.17	58.93	9Ø TH	2.32
55.88	95 TH	2.20	60.96	95 TH	2.40
57.91	99 TH	2.28	62.99	99 TH	2.48

SELLION-SUBNASALE LENGTH: The midsagittal distance between the SELLION and SUBNASALE landmarks of the nose.

TABLE 19. MIDNASAL BRIDGE HEIGHT FROM MAXILLONASALE PLANE LINE

SU	MMARY STATISTICS	3	s	MMARY STATISTICS	3.
MILLIMET	TERS	INCHES	MILLIME	TERS	INCHES
21.96	MEAN	Ø.86	25.44	MEAN	1.00
Ø.27	STD ERROR (MEAN)	Ø.Ø1	Ø.34	SID ERROR (MEAN)	0.01
2.97	SID DEVIATION	Ø. 12	3.55	SID DEVIATION	Ø. 14
13.10	MINIMIM	Ø.55	18.Ø3	MINIMIM	Ø.71
29.Ø1	MUNIXAM	1.14	39.12	HAXIMIM	1.54
COEFF.	OF VARIATION(%)	13.48	COEFF.	OF VARIATION(%)	13.94
SYMMETE	Ϋ́	Ø.Ø4	SYMMET	XX.	Ø.72
KURTOSI	S	3.11	KURTOS	rs .	4.85
NUMBER	OF SUBJECTS	119	NUMBER	OF SUBJECTS	111
	PERCENTILES			PERCENTILES	
MILLIMET	ERS	INCHES	MILLIME	TERS .	INCHES
13.97	1 51	Ø.55	18.03	ist	0.71
17.02	5 279	D. 87	29.97	HT 2	\$.79
18.Ø3	10 TH	0.71	21.08	10 TH	B. 93
20.97	25 TH	0.79	23.11	25 TH	0.91
22.10	5Ø TH	Ø.87	24.89	50 TH	Ø. 98
23.88	75 TH	Ø.94	27.94	75 TH	1.10
25.91	9Ø TH	1.Ø2	29.97	9Ø TH	1.18
26. <del>9</del> 2	95 TH	1.Ø6	29.99	95 TH	1.20
28.96	99 TH	1.14	38.10	99 TH	1.50

MIDNASALE BRIDGE HEIGHT FROM MAXILLONASALE PLANE LINE: A perpendicular projection distance of the MIDNASALE landmark away from a line coincident with the right and left MAXILLONASALE landmarks.

TABLE 20. PRONASALE HEIGHT FROM MAXILLOALARE PLANE LINE

			<del></del>		
st	MMARY STATISTICS	<b>;</b>	su	MARY STATISTIC	S
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES
32.Ø2	MEAN	1.26	37.12	MEAN	1.46
Ø. 28	STD ERROR (MEAN)	Ø.Ø1	Ø.34	STD ERROR (MEAN	Ø.Ø1
3.10	SID DEVIATION	Ø. 12	3.61	SID DEVIATION	Ø. 14
23.11	MINIMUM	Ø.91	26.92	MINIMIM	1.06
39.88	MAXIMIM	1.58	45.97	MAXIMIM	1.81
COEFF.	OF VARIATION(%)	9.55	COEFF.	OF VARIATION(%)	9.76
SYMMETE	<b>TY</b>	Ø.Ø7	SYMMETR	Y.	-Ø.36
KURTOSI	:S	3.59	KURTOSI	S	3.26
NUMBER	OF SUBJECTS	120	NUMBER	OF SUBJECTS	111
	PERCENTILES			PERCENTILES	
MILLIME	TERS	INCHES	MILLIMET	ERS	INCHES
23.11	1 ST	Ø. 91	26.92	1 ST	1.Ø6
26.92	5 TH	1.06	29.97	5 TH	1.18
27.94	10 TH	1.10	33.Ø2	10 TH	1.30
29.97	25 TH	1.18	35.Ø5		1.38
32.00	5Ø TH	1.26	37.Ø8	5Ø TH	1.46
34.04	75 TH	1.34	39.88		1.57
36.Ø7	9Ø TH	1.42	40.89	9Ø TH	1.61
37.Ø8	95 TH	1.46	42.93		1.69
39.12	99 TH	1.54	44.96	99 TH	1.77

PRONASALE HEIGHT FROM MAXILLOALARE PLANE LINE: A perpendicular projection distance of the PRONASALE landmark away from a line coincident with both bilateral MAXILLOALARE landmarks.

TABLE 21. PRONASALE HEIGHT FROM SELLION-PROMENTALE PLANE LINE

MALE

<del></del>	<del></del>		·		
st	MMARY STATISTICS	5	SU	MMARY STATISTIC	5
MILLIME	TERS	INCHES	MILLIME	ERS	INCHES
24.83	MEAN	Ø. 98	27.52	MEAN	1.09
Ø.25	STD ERROR (MEAN)	Ø.Ø1	Ø.34	SID ERROR (MEAN	Ø.34
2.95	SID DEVIATION	Ø. 12	3.62	SID DEVIATION	
19.Ø5	MINIMUM	Ø.75	16.00	MINIMIM	Ø.63
39.12	MAXIMUM	1.54	35 <b>.</b> Ø5	MAXIMIM	1.38
COEFF.	OF VARIATION(%)	11.99	COEFF.	OF VARIATION(%)	13.Ø5
SYMMETE	XX.	Ø.91	SYMMET	₹Y	-Ø.Ø3
KURTOS1	S	5.57	KURTOSI	s	2.92
NUMBER	OF SUBJECTS	144	NUMBER	OF SUBJECTS	114
	PERCENTILES			PERCENTILES	
MILLIMET	TERS	INCHES	MILLIMET	TERS	INCHES
19.05	1 ST	Ø.75	16.00	1 ST	Ø.63
20.08	5 TH	Ø.82	22.10	5 TH	Ø.87
21.Ø8	1Ø TH	Ø.83	23.11	10 TH	Ø.91
23.11	25 TH	Ø.91	24.89	25 TH	Ø.98
24.89	50 TH	Ø. 98	26.92	5Ø TH	1.26
25.91	75 TH	1.Ø2	29.97	75 TH	1.18
28.96	9Ø TH	1.14	32.00	9Ø TH	1.26
29.97	95 TH	1.18	34.04	95 TH	1.34
33.Ø2	99 TH	1.3Ø	35.Ø5	99 TH	1.38

PRONASALE HEIGHT FROM SELLION-PROMENTALE PLANE LINE: The perpendicular projection distance of PRONASALE away from a midsagittal line coincident with the SELLION and PROMENTALE landmarks.

TABLE 22. SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE

MALE

			· · · · · · · · · · · · · · · · · · ·			
SUMMARY STATISTICS		SU	Mary Statistic	TICS		
MILLIME	TERS	INCHES	MILLIMET	ĒRS	INCHES	
11.42	MEAN	Ø.45	12.87	MEAN	Ø.51	
Ø.28	STD ERROR (MEAN)	Ø.Ø2	Ø.31	SID ERROR (MEAN)	Ø.31	
3.13	STD DEVIATION	Ø. 12	3.21	STD DEVIATION	Ø. 13	
6.10	MINIMIM	Ø.24	7.00	MINIMIM	Ø.28	
22.10	MAXIMIM	Ø.87	26. <b>ØØ</b>	MAXIMIM	1.02	
COEFF.	OF VARIATION(%)	27.11	COEFF.	OF VARIATION(%)	24.85	
SYMMETE	RY	Ø.48	SYMMETR	Y	Ø.88	
KURTOS1	S	3.37	KURTOSI	S	4.93	
NUMBER	OF SUBJECTS	121	NUMBER	OF SUBJECTS	111	
	PERCENTILES			PERCENTILES		
MILLIMET	TERS	INCHES	MILLIMET	ERS	INCHES	
6.10	1 ST	Ø.24	7.Ø1	1 ST	Ø.28	
7.11	5 TH	Ø.28	8.00	5 TH	Ø.31	
7.87	10 TH	Ø.31	8.99	1Ø TH	Ø.35	
8.89	25 TH	Ø.35	11.00	25 TH	Ø.43	
1Ø. 92	5Ø TH	Ø.43	13.00	5Ø TH	Ø.51	
13.97	75 TH	Ø.55	14.00	75 TH	0.55	
14.99	9Ø TH	Ø.59	16.99	9Ø TH	Ø.67	
16.00	95 TH	Ø-63	18.Ø1	95 TH	Ø.71	
20.07	99 TH	Ø.79	23.Ø1	99 TH	Ø.91	

SELLION HEIGHT FROM MEDIAL CANTHUS PLANE LINE: A perpendicular projection distance of the SELLION landmark away from a line coincident with both bilateral areas just medial of the ENTOCANTHION landmarks.

TABLE 23. SELLION HEIGHT FROM LATERAL ORBITAL PLANE LINE

SUMMARY STATISTICS		su	MMARY STATISTICS	<b>x</b>		
MILLIMETE	irs .	INCHES	MILLIMET	ERS	INCHES	
29.56	MEAN	1.17	32.99	MEAN	1.30	
ø.5ø	STD ERROR (MEAN)	Ø.Ø1	Ø.56	SID ERROR (MEAN)	Ø.Ø2	
3.8Ø	STD DEVIATION	Ø. 15	3.44	SID DEVIATION	Ø. 13	
21.Ø8	MINIMIM	Ø.83	26.92	MINIMIM	1.05	
39.12	MAXIMIM	1.54	38.10	MAXIMUM	1.50	
COEFF. C	F VARIATION(%)	12.55	COEFF.	OF VARIATION(%)	10.14	
SYMMETRY	?	Ø.42	SYMMETR	Y	-0.14	
KURTOSIS	5	3.07	KURTOSI	S	1.77	
NUMBER C	F SUBJECTS	58 	NUMBER (	OF SUBJECTS	38	
	PERCENTILES			PERCENTILES		
MILLIMETE	<b>IRS</b>	INCHES	MILLIMET	ars	INCHES	
2Ø.83	1 ST	Ø.82	26.92	1 ST	1.Ø6	
23.88	5 TH	Ø. 94	27.94	5 TH	1.10	
24.89	10 TH	Ø.98	28.45	10 TH	1.12	
27.43	25 TH	1.Ø8	29.97	25 TH	1.18	
28.96	5Ø TH	1.14	34.Ø4	5Ø TH	1.34	
32.00	75 TH	1.26	35.Ø5	75 TH	1.38	
35.Ø5	9Ø TH	1.38	37.Ø8	9Ø TH	1.46	
36.Ø7	95 TH	1.42	37.59	95 TH	1.48	
39.12	99 TH	1.54	38.1Ø	99 TH	1.50	
				· ·		

SELLION HEIGHT FROM LATERAL ORBITAL PLANE LINE: A perpendicular projection distance of the SELLION landmark away from a line coincident with the antero-lateral aspects of the lateral orbital rims, just lateral of the right and left ECTOCANTHION landmarks.

## 

# 

DATE: 4-93